



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

and on walking to B , a distance a , directly down the hill towards the foot of the pole the white part subtends the same angle. What is the length of the white part, if the point B is at a distance b from the foot of the pole?

Solution by G. B. M. ZERE, A. M., Ph. D., Professor of Mathematics and Applied Science, Texarkana College, Texarkana, Arkansas-Texas.

Let DE be the length painted white; then a circle will pass through A , B , D , E . Let $\angle EAD = \angle EBD = \alpha$, $AB = a$, $BC = b$, $\angle DAB = \angle DEB = \theta$, $\angle ABE = \angle ADE = \varphi$, $DC = y$, and $DE = x$.

$$\text{Then } (x+y)y = (a+b)b \dots \dots \dots (1).$$

$$AE : a = \sin \varphi : \sin(\alpha + \theta + \varphi), \quad x : AE = \sin \alpha : \sin \varphi.$$

$$\therefore x = \frac{a \sin \alpha}{\sin(\alpha + \theta + \varphi)} \dots \dots \dots (2),$$

$$b : x + y = \sin \theta : \sin(\alpha + \varphi) \dots \dots \dots (3),$$

$$(x+y) : a+b = \sin(\alpha + \theta) : \sin(\alpha + \varphi) \dots \dots \dots (4).$$

Eliminating θ between (3) and (4),

$$\left\{ \frac{(x+y)^4}{(a+b)^2} - \frac{2b(x+y)^2 \cos \alpha}{a+b} + b^2 \right\} \sin^2(\alpha + \varphi) = (x+y)^2 \sin^2 \alpha \dots \dots \dots (5).$$

Eliminating θ between (2) and (3),

$$\begin{aligned} & [\{b^2 x^2 - x^2(x+y)^2\}^2 + 4a^2 b^2 x^2(x+y)^2 \sin^2 \alpha] \sin^4(\alpha + \varphi) \\ & - 2a^2 \sin^2 \alpha (x+y)^2 \{b^2 x^2 + x^2(x+y)^2\} \\ & \sin^2(\alpha + \varphi) + a^4(x+y)^4 \sin^4 \alpha = 0 \dots \dots \dots (6). \end{aligned}$$

Eliminating $\sin(\alpha + \varphi)$ between (5) and (6) we get an equation in x and y which with (1) gives us the value of x .

Solved with result in terms of EC by *A. H. HOLMES*, and *FREDERICK R. HONEY*.


PROBLEMS.

58. Proposed by *I. J. SCHWATT*, Ph. D., Instructor in Mathematics, University of Pennsylvania, Philadelphia, Pennsylvania.

1. The point of intersection K_a' of the tangent drawn to the circumcircle about the triangle ABC at A and the side BC is harmonic conjugate to K_a with respect to BC . (K_a is the point where the symmedian line through A of the triangle ABC meets the side BC .)

2. The point K_a' is the center of the Apollonius circle passing through A of the triangle ABC .

3. Grebes point is on the line joining the middle point of any side of a triangle with the middle point of the altitude to this side.



59. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Show that the tangent plane at any point of the surface $a^2x^2 + b^2y^2 + c^2z^2 = 2bcyz + 2acxz + 2abxy$ intersects the surface $ayz + bzx + cxy = 0$ in two straight lines at right angles to one another.

CALCULUS.

Conducted by J. M. COLAW, Monterey, Va. All contributions to this department should be sent to him.

SOLUTIONS OF PROBLEMS.

47. Proposed by Prof. J. SCHEFFER, A. M., Hagerstown, Maryland.


The floor of a vault forms a square, and all sections parallel to it are squares. The two vertical sections through the middle points of the opposite sides of the floor are equal semi-circles. Find the convex surface and the volume of the vault.

I. Solution by C. W. M. BLACK, Professor of Mathematics in Wesleyan Academy, Wilbraham, Massachusetts; O. W. ANTHONY, M. Sc., Professor of Mathematics in New Windsor College, New Windsor, Maryland; A. H. HOLMES, Brunswick, Maine, and the PROPOSER.

Let $ABCD$ represent the base square, side $= 2a$, and KEI and GFH the two equal semi-circles, radius $= a$. Let $LMNO$ be another square parallel to the base square, and at the distance $PE = x$ from it. The area of $LMNO$ is $= 4(a^2 - x^2)$,

$$\therefore \text{Vol.} = 4 \int_0^a (a^2 - x^2) dx = \frac{8}{3} a^3.$$

Denoting $\angle PEQ$ by θ , we have for the surface



$$\int_0^{\frac{1}{2}\pi} 8a \cos \theta d(a\theta) = 8a^2. \quad \text{Or for the volume, } dV = 4a^2 \cos^2 \theta dx, \text{ where } x \text{ is}$$

the vertical distance, $x = a \sin \theta$; $dx = a \cos \theta d\theta$.

$$\therefore V = 4a^3 \int_0^{\frac{1}{2}\pi} \cos^3 \theta d\theta = a^3 \int_0^{\frac{1}{2}\pi} (\cos^3 \theta + 3\cos \theta) d\theta = \frac{8a^3}{3}.$$

II. Solution by BENJ. F. YANNEY, A. M., Professor of Mathematics in Mount Union College, Alliance, Ohio.

The convex surface of the vault is equivalent to the surface of a right cir-